Medical Imaging Centre

ALL OF YOUR IMAGING NEEDS

BRING OHIP CARD		CLINICAL INFORMATION	
Patient Name: Patient Tel:			
Health Card No.:			
Referring MD:		Billing #: Physician	
Appointment Time:		Signature:	
NUCLEAR IMAGING	GENERAL ULTRASOUND	OBSTETRICA	L ULTRASOUND
 Exercise Stress Persantine Stress Rest Thallium Rest Muga 	 Abodomen Abdomen + Pelvis Inguinal Female Pelvis Male Pelvis With Kidneys 	 Dating Problem Nuchal Lucency IPS Anatomy 	FOLLOW UP O Head O Face O Heart O Spine O Kidneys
VASCULAR IMAGING	O Transrectal Prostate	(11-18 Weeks)	⊖ Growth
 Carotid Doppler Arterial Doppler Upper (R) (L)(Both) OLower (R) (L)(Both) AAA Screen (limited abdominal + pelvic doppler) Venous Doppler Upper (R) (L)(Both) OLower (R) (L)(Both) Renal Artery Doppler 	 Scrotum / Testes Head & Neck Thyroid Parotid 	🔿 врр	 Placenta Amniotic Fluid Position
	○ Submandibular	BONE MINERAL DENSITY	
	 Lymph nodes Other 	 Routine (Every 3 Years) High Risk (Every 12 Months) 	
PLEASE CIRCLE/MARK REQUIRED TESTS			
istory istory			

facility providing services required



905.881.1455

GENERAL INFORMATION

BRING YOUR OHIP TO EVERY APPOINTMENT BRING A LIST OF ALL YOUR MEDICATIONS

NUCLEAR IMAGING PREPARATION

- Wear comfortable, loose clothing and walking shoes.
- Breakfast consisting of toast, jam, juice and water ONE HOUR before your test.
- 24 Hours No caffeine, No Coffee, No Decaffeinated products, No Medication with caffeine.
- Please inform us if you are on medications such as: Viagra, Cialis, Levitra etc... Do not take for 48 hrs before test
- The test will take approximately 4 hours

ABDOMEN:

AM appointments - Fat free dinner night before. Nothing to eat or drink after midnight PM appointments - Fat free breakfast before 9 am then clear fluids (no milk products). Nothing 2 hours prior to study.

ABDOMEN & PELVIC TOGETHER:

AM appointments - Nothing to eat after midnight drink 40 oz - of water (5 eight oz. glass or 1.5L). Must finish drinking ONE hour before study. DO NOT void before the test.

PM appointments - Eat fat free breakfast Must finish drinking 40 oz. of fluid ONE hour before study. DO NOT void before the test.

ALL PELVIS: Female, Male or obstetrics (less than 22 weeks)


Drink 40 oz. of water (5 eight oz. glass or 1.5L). Must finish drinking ONE hour before study. Do not void before the test Bladder must be full or test may have to be rebooked.

PREGNANCY: Greater than 22 weeks

Drink 35 oz. (1L) of water Must finish drinking ONE hour before study. Do not void before the test. Bladder must be full, or test may be rebooked.

MALE PROSTATE: Transrectal

Purchase Fleet Enema from pharmacy: follow all instructions. Proceed with enema 2 hour prior to exam. Drink 40 oz. of water Finish drinking ONE hour before study. Do not void before study.



Thornhill

398 Steeles Ave. West Suite 11-14, L4J 6X3 Tel. 905. 881. 1455 Fax. 905. 881. 1721 Central Booking Line: 416. 461. 9954

This requisition form can be taken to any licensed facility providing services required